

THE SCHOTTKY PROBLEM IN GENUS FIVE

CHARLES SIEGEL

ABSTRACT. In this paper, we present a solution to the Schottky problem in the spirit of Schottky and Jung for genus five curves. To do so, we exploit natural incidence structures on the fibers of several maps to reduce all questions to statements about the Prym map for genus six curves. This allows us to find all components of the big Schottky locus and thus, to show that the small Schottky locus introduced by Donagi is irreducible.

CONTENTS

Introduction	1
1. Background	3
2. Line Configurations	10
3. Degenerations of Abelian varieties	15
4. Contracted loci	17
5. Schottky-Jung locus	18
References	18

INTRODUCTION

The Schottky problem has a long history and is one of the first questions to be asked in complex curve theory: how can we characterize Jacobians among all abelian varieties? The question turns out to be quite subtle, and a completely satisfactory answer is still lacking, despite several characterizations existing and being a fertile ground for new techniques in studying the moduli of abelian varieties.

Historical Overview. The oldest approach to the Schottky problem is via theta functions. Certain theta functions give natural coordinates on $\mathcal{A}_g$, called thetanulls, and Schottky [Sch88] for genus four and Schottky and Jung [SJ09] for genus $g \geq 5$ conjectured equations for the locus of Jacobians. We denote the locus these equations cut out by $\mathcal{S}_g$, the Schottky locus, and then the Schottky-Jung conjecture is that $\mathcal{S}_g = \mathcal{J}_g$. Schottky and Jung's results implied only $\mathcal{J}_g \subset \mathcal{S}_g$, though that $\mathcal{J}_g$ is an irreducible component of $\mathcal{S}_g$ was shown by [Gee84]. A solution in the genus four case was announced in the late 1960s by Igusa [Igu69] and appeared independently in [Igu81] and [Fre83]. The method was to prove that $\mathcal{S}_4$ was an irreducible divisor on $\mathcal{A}_4$, and thus, must be identical with $\mathcal{J}_4$, an irreducible divisor contained inside it. Unfortunately, this method cannot be employed directly in higher genus, as it relied on the fact that $\mathcal{S}_4$ is given by a single equation in theta nulls.

Another attempt to describe the Jacobians involves the dimension of the singular locus of the theta divisor. The Riemann singularity theorem implies that this

singular locus has dimension at least $g - 4$, and this is a special condition on abelian varieties. Andreotti and Mayer [AM67] studied the locus $\mathcal{AM}_g$ of such abelian varieties and showed that it gives a weak solution to the Schottky problem, that is, the Jacobians are an irreducible component of $\mathcal{AM}_g$, but other components exist. In genus four [Bea77a] and five [Don81, Deb90] the locus $\mathcal{AM}_g$ has been completely described as

$$\mathcal{AM}_4 = \theta_{null,4} \cup \mathcal{J}_4 \text{ and } \mathcal{AM}_5 = \mathcal{A}_1 \times \mathcal{A}_4 \cup \mathcal{BE} \cup \mathcal{J}_5,$$

where $\mathcal{BE}$ is the locus of bielliptic Prym varieties and $\theta_{null,g}$ the locus of abelian varieties with a vanishing thetanull, with all the non-Jacobians contained in $\theta_{null,5}$. This approach has not been carried out in higher genus fully, but singularities of theta divisors remains an active area of study [CM08, CG00, GSM07].

A third approach has been through trisecants. Fay's trisecant formula [Fay73] tells us that the natural embedding of $\mathcal{J}(C)/\pm 1 \rightarrow \mathbb{P}^{2^g-1}$ has a four dimensional family of trisecant lines. Welters [Wel84] proved that if $A/\pm 1 \rightarrow \mathbb{P}^{2^g-1}$ has a one parameter family of trisecant lines, then it must be a Jacobian. The family of trisecants has an infinitesimal formulation as the KP hierarchy, and Novikov conjectured that an abelian variety is a Jacobian if and only if the corresponding theta function satisfies the KP hierarchy. This was proved by Shiota [Shi86]. The strongest result on trisecants is the recent proof, using difference equations, by Krichever [Kri10] that a single trisecant on $A/\pm 1$ suffices to determine that an abelian variety is a Jacobian.

This paper. In [Don87a, Don87b], Donagi showed that the original Schottky-Jung conjecture was incorrect. In particular, he showed that $\mathcal{C}$, the locus of intermediate Jacobians of cubic threefolds, is contained in $\mathcal{S}_5$. Additionally, he offered a means of correcting this, and conjectured that nothing not accounted for by his methods appears in the Schottky locus. However, even in genus five a proof was out of reach at the time, with some aspects of degenerations of dimension five abelian varieties remaining unclear until Izadi's thesis [Iza91].

In this paper, we present a proof of

Theorem. *Inside of $\mathcal{A}_g$, we have*

$$\mathcal{J}_5 = \mathcal{S}_g^{\text{small}}.$$

The paper is organized as follows. In section 1, we will gather results from the literature, recall definitions and set notation for the remainder of the paper. Section 2 consists of the geometry and combinatorics of point and line configurations over $\mathbb{F}_2$, and describes incidence structures on the fibers of several maps defined below. Section 3 studies degenerations of abelian varieties of dimension five, using results from Izadi's thesis [Iza91] to complete the computation of the degree of the map β_5 begun in section 2 via incidence structures. Section 4 consists of a study of the contracted loci for β_5 , completing our picture of this map and allowing us to proceed to section 5, where we use this to compute the Schottky locus, and prove the main theorem.

Acknowledgments. I would like to thank first of all Ron Donagi, who suggested this problem to me and advised me through the work as my Ph.D. thesis at the University of Pennsylvania. Also, I am indebted to Sam Grushevsky, Gavril Farkas, Angela Ortega, Tony Pantev, Elham Izadi and Angela Gibney for conversations

clarifying aspects of the problem as well as the geometry of the moduli spaces involved in this paper. This work was supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, and also by the University of Pennsylvania

1. BACKGROUND

1.1. Moduli of curves and abelian varieties. First, we will describe the various moduli spaces that we will need, all of which are covers of the spaces $\mathcal{M}_g$ and $\mathcal{A}_g$, the moduli space of genus g compact Riemann surfaces and of g dimensional principally polarized abelian varieties, respectively.

We will need first $\mathbb{H}_g$, the Siegel upper half space of $g \times g$ symmetric complex matrices with positive definite imaginary part. This space is the stack universal cover of $\mathcal{A}_g$, and has an action of $\Gamma_g = \mathrm{Sp}(2g, \mathbb{Z})$ by

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \cdot \Omega = (A\Omega + B)(C\Omega + D)^{-1}$$

such that $\mathcal{A}_g \cong \mathbb{H}_g / \Gamma_g$ as complex analytic spaces.

Other space we will need are defined by subgroups of Γ_g . We set $\mathcal{A}_g(n)$ to be the quotient of $\mathbb{H}_g$ by

$$\Gamma_g(n) = \{\gamma \in \Gamma_g \mid \gamma \equiv 1 \pmod{n}\}.$$

This space parameterizes pairs (A, ϕ) where $A \in \mathcal{A}_g$ and $\phi: A[n] \rightarrow (\mathbb{Z}/n\mathbb{Z})^{2g}$ is a symplectic isomorphism from the points of order n on A to $(\mathbb{Z}/n\mathbb{Z})^{2g}$. For theta functions, it will be essential to use $\mathcal{A}_g(n, 2n)$, which is the quotient by

$$\Gamma_g(n, 2n) = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g(n) \mid \mathrm{diag}({}^t AC) \equiv \mathrm{diag}({}^t BD) \equiv 0 \pmod{2n} \right\}.$$

For the Prym map, we will need to fix the vector $v = (0, \dots, 0, 1/2) \in \mathbb{Q}^{2g}$, and then look at the groups

$$\begin{aligned} \mathcal{R}\Gamma_g(n) &= \{\gamma \in \Gamma_g(n) \mid \gamma \cdot v \equiv v \pmod{n\mathbb{Z}^{2g}}\} \\ \mathcal{R}\Gamma_g(n, 2n) &= \{\gamma \in \Gamma_g(n, 2n) \mid \gamma \cdot v \equiv v \pmod{n\mathbb{Z}^{2g}}\} \end{aligned}$$

corresponding to the moduli spaces $\mathcal{RA}_g(n)$ and $\mathcal{RA}_g(n, 2n)$. These spaces parameterize pairs (A, μ) where A is in $\mathcal{A}_g(n)$ or $\mathcal{A}_g(n, 2n)$, respectively, along with $\mu \in A[2]$ nonzero.

For $\mathcal{M}_g$, we will need all of the analogous spaces to the ones defined above for $\mathcal{A}_g$. They can all be constructed via pullbacks: we use the Torelli map $\mathcal{J}_g: \mathcal{M}_g \rightarrow \mathcal{A}_g$ and take the fiber product with the projection to $\mathcal{A}_g$. For example, $\mathcal{RM}_g$ is the space of pairs (C, μ) where μ is a nonzero point of order 2 on $\mathcal{J}(C)$. We will need one additional space, $\mathcal{R}^2\mathcal{M}_g$, whose elements are ordered triples (C, μ, ν) with $C \in \mathcal{M}_g$, μ, ν distinct nonzero points of order 2 on the Jacobian such that for any line bundle L on C with $L^{\otimes 2} \cong K_C$, we have

$$(1) \quad h^0(C, L) + h^0(C, L \otimes \mu) + h^0(C, L \otimes \nu) + h^0(C, L \otimes \mu \otimes \nu) \equiv 0 \pmod{2}.$$

The left hand side is called the *Weil pairing* and we will refer to pairs satisfying 1 as orthogonal. For abelian varieties that are not Jacobians, we can define the Weil pairing as the intersection pairing on $H_1(A, \mathbb{Z}/2\mathbb{Z})$. (That these are the same is shown in [Mum71])

Finally, here we note another interpretation of $\overline{\mathcal{RM}}_g$. The data of a smooth curve C and $\mu \in \mathcal{J}(C)[2]$ is equivalent to $\tilde{C} \rightarrow C$, an unramified double cover. We can construct $\tilde{C}$ inside the total space of μ by fixing $\varphi : \mu^{\otimes 2} \rightarrow \mathcal{O}_C$ an isomorphism and then, on sufficiently small open sets $U \subset C$, we have

$$\tilde{C}|_U = \{\sigma^2 = 1 | \sigma \in \mu(U)\}.$$

We will use (C, μ) and $(C, \tilde{C})$ interchangeably, along with $(\tilde{C}, \iota)$ where $\iota : \tilde{C} \rightarrow \tilde{C}$ is the involution corresponding to the double cover.

1.2. Compactifications. In this paper, we will need several partial compactifications of the moduli spaces we study, particularly $\mathcal{M}_g$, $\mathcal{A}_g$, $\mathcal{RA}_g$, $\mathcal{RA}_g$ and $\mathcal{R}^2\mathcal{M}_g$.

We will start with the boundary for $\mathcal{A}_g$ and $\mathcal{RA}_g$, which is simpler than that of $\mathcal{M}_g$ and its covers. The space $\mathcal{A}_g$ has a distinguished compactification, the Satake compactification, which all other compactifications map to. We will only need the corank 1 part, which we will denote by $\overline{\mathcal{A}}_g^s$ and we note that $\partial\overline{\mathcal{A}}_g^s \cong \mathcal{A}_{g-1}$. There is a larger class of compactifications called toroidal compactifications [AMRT10]. We will only need the corank 1 part here as well, and for all toroidal compactifications, this is the same, so we write $\overline{\mathcal{A}}_g^t = \text{Bl}_{\mathcal{A}_{g-1}} \overline{\mathcal{A}}_g^s$. This has the property that $\partial\overline{\mathcal{A}}_g^t$ is a divisor, and is isomorphic to $\mathcal{X}_{g-1}$, the universal Kummer variety over $\mathcal{A}_{g-1}$. We will interpret it as parameterizing $\mathbb{C}^\times$ -extensions of abelian varieties of dimension $g-1$.

The space $\mathcal{RA}_g$ has a slightly more complex boundary, and for reference we reproduce results in [Don87a] describing it. There are both Satake and toroidal compactifications, for which we will again only need the corank 1 part. In either case, there are three components, depending on the relationship between the point of order two, μ , and the vanishing cycle δ taken modulo two:

- $\partial^I \overline{\mathcal{RA}}_g^s \cong \mathcal{A}_{g-1}$ and $\partial^I \overline{\mathcal{RA}}_g^t \cong \mathcal{X}_{g-1}$ are the components where $\delta = \mu$.
- $\partial^{II} \overline{\mathcal{RA}}_g^s \cong \mathcal{RA}_{g-1}$ and $\partial^{II} \overline{\mathcal{RA}}_g^t \cong \mathcal{RX}_{g-1}$ (The fiber of $\mathcal{RX}_{g-1} \rightarrow \mathcal{RA}_{g-1}$ over $(X, \mu) \in \mathcal{RA}_{g-1}$ is the double cover of X determined by $(\mu)^\perp$) are the components where $\delta \neq \mu$ and the two points are orthogonal.
- $\partial^{III} \overline{\mathcal{RA}}_g^s \cong \mathcal{A}_{g-1}$ and $\partial^{III} \overline{\mathcal{RA}}_g^t \cong \mathcal{X}_{g-1}$ are the components where δ and μ are not orthogonal.

In fact, we have

Proposition 1.1 ([Don87a, Proposition 2.3.2]).

The projection $\overline{\mathcal{RA}}_g^t \rightarrow \overline{\mathcal{RA}}_g^s$ is simply ramified along $\partial^{III} \overline{\mathcal{RA}}_g^t$ and unramified on the other boundary components. (Note that the map $\partial^{III} \overline{\mathcal{RA}}_g^t \rightarrow \partial^{III} \overline{\mathcal{RA}}_g^s$ is given by $\mathcal{RX}_{g-1} \xrightarrow{\times 2} \mathcal{RX}_{g-1} \rightarrow \mathcal{RA}_{g-1}$ with the first map multiplication by two along the fibers)

For $\mathcal{M}_g$ and its covers, the boundary is somewhat more complex. We will only be using partial compactifications contained in the Deligne-Mumford stable curve compactification [DM69]. In this compactification, we have δ_i for $1 \leq i \leq \lfloor \frac{g-1}{2} \rfloor$ consisting of reducible curves with a component of genus i meeting a component of genus $g-i$ at a point. There is also a component δ_0 consisting of irreducible curves of geometric genus $g-1$ with a single node [HM98].

The situation for $\overline{\mathcal{RM}}_g$ is slightly more complex. Each boundary component of $\overline{\mathcal{M}}_g$ splits into three components. The reducible components δ_i give us ∂_i , ∂_{g-i}

and $\partial_{i,g-i}$, consisting of pairs (C, μ) where μ is supported on the curve of genus i , $g-i$ or both. For δ_0 , we have ∂^I , ∂^{II} and ∂^{III} as with $\overline{\mathcal{RA}}_g^s$. We also, for later reference, describe the corresponding double covers:

Proposition 1.2 ([Don92, Examples 1.9]).

The generic element of the boundary components of $\overline{\mathcal{RM}}_g$ correspond to double covers as follows:

- ∂_i for $1 \leq i \leq g-1$: the base curve is $C = C_i \cup_p C_{g-i}$ and the double cover is $\tilde{C} = C_{g-i} \cup_{p_1} \tilde{C}_i \cup_{p_2} C_{g-i}$.
- $\partial_{i,g-i}$ for $1 \leq \lfloor \frac{g-1}{2} \rfloor$: the base curve is $C = C_i \cup_{p \sim q} C_{g-i}$ and $\tilde{C} = \tilde{C}_{g-i} \cup_{p_1 \sim q_1, p_2 \sim q_2} \tilde{C}_i$.
- ∂^I : the base curve is $C = X/p \sim q$ and the double cover is $\tilde{C} = X_0 \coprod X_1/p_0 \sim q_1, p_1 \sim q_0$. This is called a Wirtinger double cover.
- ∂^{II} : the base curve is $C = X/p \sim q$ and the double cover is $\tilde{C} = \tilde{X}/p_0 \sim q_0, p_1 \sim q_1$ where $\tilde{X} \rightarrow X$ is an unramified double cover. This is called an unallowable double cover.
- ∂^{III} : the base curve is $C = X/p \sim q$ and the double cover is $\tilde{C} = \tilde{X}/\tilde{p} \sim \tilde{q}$ where $\tilde{X} \rightarrow X$ is a double cover ramified over p and q . This is called a Beauville double cover.

1.3. The Prym map. In this section, we will define, describe and collect useful results about the Prym map.

Let C be a genus g curve and $\pi: \tilde{C} \rightarrow C$ be any morphism of curves. Then we can define a map $\text{Nm}_\pi: \mathcal{J}(\tilde{C}) \rightarrow \mathcal{J}(C)$ by writing $D \in \mathcal{J}(\tilde{C})$ as $D = \sum n_P P$ and setting $\text{Nm}_\pi(D) = \sum n_P \pi(P)$. If π is surjective, then so is Nm_π . This is especially useful in the case where π is an unramified double cover. In this case, the kernel is $g-1$ dimensional and has two components. We define $\mathcal{P}_g(C, \tilde{C}) = \ker^0 \text{Nm}_\pi$, the connected component of the identity of the kernel of the norm map, and call it the Prym variety of the double cover. The principal polarization on $\mathcal{J}(\tilde{C})$ restricts to twice a principal polarization on $\mathcal{P}(C, \tilde{C})$ [Mum74, Corollary 2], thus giving $\mathcal{P}(C, \tilde{C})$ the natural structure of a principally polarized abelian variety, giving us a map $\mathcal{P}_g: \mathcal{RM}_g \rightarrow \mathcal{A}_{g-1}$. Additionally, we have:

Lemma 1.3 (Mumford Sequence [Mum74, Corollary 1]).

Let $(C, \mu) \in \mathcal{RM}_g$. Then we have a short exact sequence

$$0 \rightarrow \langle \mu \rangle \rightarrow \mu^\perp \rightarrow \mathcal{P}(C, \mu)[2] \rightarrow 0$$

where we are taking the orthogonal complement with respect to the Weil pairing.

If $\nu \in \mu^\perp$ we will denote the image of ν in $\mathcal{P}(C, \mu)[2]$ by $\bar{\nu}$.

This allows us to define also $\mathcal{RP}_g: \mathcal{R}^2\mathcal{M}_g \rightarrow \mathcal{RA}_{g-1}$ by $\mathcal{RP}_g(C, \mu, \nu) = (\mathcal{P}(C, \mu), \bar{\nu})$.

Unfortunately, as defined, the Prym map is not proper, so we must study the Prym map on the boundary of $\mathcal{RM}_g$.

Proposition 1.4 ([Don92, Example 1.9]).

The Prym map extends to the general point of the boundary of $\overline{\mathcal{RM}}_g$ as follows along each component:

- ∂^I : For a Wirtinger double cover, $\mathcal{P}_g(C, \tilde{C}) \cong \mathcal{J}_{g-1}(X)$.

- ∂^{II} : For an unallowable double cover, $\mathcal{P}_g(C, \tilde{C})$ is the $\mathbb{C}^\times$ -extension of $\mathcal{P}_{g-1}(X, \tilde{X})$ given by $p_0 - p_1 + q_0 - q_1$ in $\mathcal{P}_{g-1}(X, \tilde{X})$
- ∂^{III} : The Prym of a Beauville double cover is a principally polarized abelian variety, and this and other ramified Prym maps are studied in [MP12]. It is isogenous to $\mathcal{P}_{g-1}(X, \tilde{X})$ but is not isomorphic to it.
- ∂_i fits into a diagram:

$$\begin{array}{ccc} \partial_i & \xrightarrow{\mathcal{P}_g} & \mathcal{A}_{i-1} \times \mathcal{A}_{g-i} \\ \downarrow & \nearrow \mathcal{P}_i \times \mathcal{J}_{g-i} & \\ \mathcal{RM}_i \times \mathcal{M}_{g-i} & & \end{array}$$

- $\partial_{i,g-i}$ fits into a diagram:

$$\begin{array}{ccc} \partial_{i,g-i} & \xrightarrow{\mathcal{P}_g} & \partial \overline{\mathcal{A}}_{g-1}^t \\ \downarrow & & \downarrow \\ \mathcal{RM}_i \times \mathcal{RM}_{g-i} & \xrightarrow{\mathcal{P}_i \times \mathcal{P}_{g-i}} & \mathcal{A}_{k-1} \times \mathcal{A}_{g-k-1} \end{array}$$

Unfortunately, $\mathcal{P}_g$ is only a rational map, it cannot be extended to all over $\overline{\mathcal{RM}}_g$. However, if we look at $\mathcal{P}_g^{-1}(\mathcal{A}_{g-1}) = (\overline{\mathcal{RM}}_g)^{\text{allowable}}$, then we have

Theorem 1.5 (Prym is Proper [DS81, Theorem 1.1], [Bea77a, Proposition 6.3]). *For all g , $(\overline{\mathcal{RM}}_g)^{\text{allowable}} \rightarrow \mathcal{A}_{g-1}$ is proper and for $g \leq 6$ it is surjective.*

Finally, we will need a theorem of Shokurov, based on Mumford's analysis of the singularities of theta divisors:

Proposition 1.6 ([Sho82], stated as it appears in [Bea89]).

Assume $g \neq 4$. If $\mathcal{P}_g(C, \tilde{C})$ is a Jacobian, then C is hyperelliptic, trigonal or is a plane quintic with an even point of order two.

In particular, this theorem implies that C is not bielliptic, that is, it is not a branched cover of an elliptic curve, and this will be the main use we will have for it later.

1.4. Fibers of the Prym map. Any study of the fibers of the Prym map begins with the tetragonal construction of Donagi, announced in [Don81] and more fully developed in [Don92]. Let $\pi: \tilde{C} \rightarrow C$ be an unramified double cover and $f: C \rightarrow \mathbb{P}^1$ be a degree four map. Then we define

$$f_*\tilde{C} = \{D \in \text{Sym}^4 \tilde{C} \mid \text{Nm}_\pi(D) = f^{-1}(k) \text{ for some } k \in \mathbb{P}^1\}.$$

The involution ι on $\tilde{C}$ extends to $f_*\tilde{C}$, which also splits into two components, giving us two new towers $\tilde{C}_0 \xrightarrow{\pi_0} C_0 \xrightarrow{f_0} \mathbb{P}^1$ and $\tilde{C}_1 \xrightarrow{\pi_1} C_1 \xrightarrow{f_1} \mathbb{P}^1$ where $(C_i, \tilde{C}_i) \in \overline{\mathcal{RM}}_g$ and f_i is of degree four. The key theorem is

Theorem 1.7 ([Don81, Proposition 1.1]).

The tetragonal construction commutes with the Prym map:

$$\mathcal{P}_g(C, \tilde{C}) \cong \mathcal{P}_g(C_0, \tilde{C}_0) \cong \mathcal{P}_g(C_1, \tilde{C}_1).$$

This is a generalization of a theorem of Recillas identifying trigonal Pryms:

Theorem 1.8 (The Trigonal Construction [Don92, Example 2.15(1)], [Rec74]).
If $C \rightarrow \mathbb{P}^1$ is tetragonal and $\tilde{C} \rightarrow C$ is the reducible double cover, then the tetragonal construction gives a copy of $\tilde{C} \rightarrow C \rightarrow \mathbb{P}^1$ and also $\mathbb{P}^1 \cup \tilde{T} \cup \mathbb{P}^1 \rightarrow \mathbb{P}^1 \cup T \rightarrow \mathbb{P}^1$ where T is a trigonal curve and $\mathcal{P}_g(T, \tilde{T}) \cong \mathcal{J}_{g-1}(C)$. Additionally, this operation is a bijection, so every trigonal Prym is a tetragonal Jacobian.

and also we recall a theorem of Mumford's identifying hyperelliptic Pryms:

Proposition 1.9 ([Mum71]).

The Pryms of hyperelliptic curves are products of Jacobians.

Now we will discuss the fibers of the Prym maps $\mathcal{P}_6$ and $\mathcal{P}_5$, as well as the behavior over certain special loci. For a general genus six curve, there are exactly five g_4^1 's, and from this and the fact that any two points in the fiber of $\mathcal{P}_6$ are related by at most two applications of the tetragonal construction, we get

Theorem 1.10 ([DS81, Theorem 2.1][Don81, Theorem 2.1]).

The map $\mathcal{RM}_6 \rightarrow \mathcal{A}_5$ is generically finite of degree 27 with Galois group WE_6 , the Weil group of the root system of type E_6 , giving the general fiber the structure of the 27 lines on a cubic surface.

We can see this structure clearly over some special loci in $\mathcal{A}_5$. However, first we need to perform some blowups. Let $\mathcal{RQ}^0$ and $\mathcal{RQ}^1$ be the loci of plane quintic curves with an odd or even point of order two, respectively. (If μ is a point of order two on $\mathcal{J}(Q)$, its parity is the parity of the dimension of the space of global sections of $\mu \otimes g_5^2$.) Also let $\mathcal{RM}_6^{\text{Trig}}$ be the locus of genus six curves with a g_3^1 , let $\mathcal{C}$ be the locus of intermediate Jacobians of cubic threefolds in $\mathcal{A}_5$. Finally, set $\widetilde{\mathcal{RM}}_6$ to be the blowup of $\mathcal{RM}_6$ along $\mathcal{RQ}^0 \cup \mathcal{RQ}^1 \cup \mathcal{RM}_6^{\text{Trig}}$ and $\tilde{\mathcal{A}}_5$ the blowup of $\mathcal{A}_5$ along $\mathcal{J}_5 \cup \mathcal{C}$. Then the Prym map $\tilde{\mathcal{P}}_6 : \widetilde{\mathcal{RM}}_6 \rightarrow \tilde{\mathcal{A}}_5$. This blown up Prym map is easier to analyze and gives more explicitly the structure of the tetragonal construction and the fibers.

Proposition 1.11 ([Don92, Remark 4.5.1]).

Any two points in a fiber of $\tilde{\mathcal{P}}_6$ are related by at most two tetragonal constructions.

Proposition 1.12 ([Don92, 4.3.4]).

The fiber of $\mathcal{P}_6$ over $X \in \mathcal{C}$ is the Fano surface of lines in X . In $\tilde{\mathcal{A}}_5$, the inverse image of $X \in \mathcal{C}$ is pairs (X, H) where H is a hyperplane in $\mathbb{P}^4$. Then, $\tilde{\mathcal{P}}_6^{-1}(X, H) = \{\ell \mid \ell \subset X \cap H\}$.

The above proposition exhibits the structure of the 27 lines very explicitly, and it follows from [Bea77b, Theorem 2.1] which uses the correspondence between conic bundle structures and lines on a cubic threefold to identify the intermediate Jacobians with the Pryms of discriminant curves.

Suppressing normal data, we can also describe the fiber over the Jacobian of a curve:

Proposition 1.13 ([Don92, 4.3.7]).

Let $C \in \mathcal{M}_5$. Then $\tilde{\mathcal{P}}_6^{-1}(\mathcal{J}(C))$ contains one plane quintic with an odd point of order two, ten trigonal curves with double covers and sixteen Wirtinger double covers.

Another extremely useful result is

Proposition 1.14 ([Don81, Corollary 2.3],[Don92, 4.8],[Cle83]).

The ramification locus of $\mathcal{P}_6$ is mapped six to one onto the branch locus, which is the locus of intermediate Jacobians of quartic double solids.

Finally, for $\mathcal{P}_6$ we look over the boundary where we have

Proposition 1.15 ([Iza91, Corollary 6.5]).

The restriction of the Prym map to $\partial^{II} \rightarrow \partial \overline{\mathcal{A}}_5^t$ is generically unramified and finite.

Now, we recall the major results on $\mathcal{P}_5$:

Proposition 1.16 ([Don92, Theorem 5.2]).

- (1) *There exists an involution $\lambda : \mathcal{RM}_5 \rightarrow \mathcal{RM}_5$ such that $\mathcal{P}_5 \circ \lambda = \mathcal{P}_5$.*
- (2) *There is a natural birational map $\chi : \overline{\mathcal{A}}_4^t \dashrightarrow \overline{\mathcal{RC}}^0$, where $\overline{\mathcal{RC}}^0$ is the space of cubic threefolds with an even point of order two on the intermediate Jacobian, including nodal cubic threefolds as limits.*

Given these two maps, Donagi showed

Proposition 1.17 ([Don81, Theorem 3.3]).

For $A \in \mathcal{A}_4$ generic, $\mathcal{P}_5^{-1}(A)/\lambda$ is isomorphic to the Fano surface of lines in $\chi(A)$, and $(C_1, \tilde{C}_1)$ and $(C_2, \tilde{C}_2)$ are tetragonally related if and only if the corresponding lines intersect. Thus, any two points in $\mathcal{P}_5^{-1}(A)$ are related by at most two tetragonal constructions.

Before moving on, we will identify the fibers of $\mathcal{P}_5$ over Jacobians:

Proposition 1.18 ([Don92, Theorem 5.14]).

Let $B \in \mathcal{M}_4$ be a general curve of genus 4 and let $(X, \delta) = \chi(\mathcal{J}(B))$. Then

- (1) *X is a nodal cubic threefold*
- (2) *The double cover of the Fano surface of lines in X is reducible and each component is isomorphic to $\text{Sym}^2 B$*
- (3) *$\mathcal{P}_5^{-1}(\mathcal{J}(B))$ is isomorphic to the double cover of the Fano surface of lines of X , with one component of trigonal curves $T_{p,q}$ and one of Wirtinger double covers $S_{p,q}$, for $(p, q) \in \text{Sym}^2 B$*
- (4) *The tetragonal construction takes $S_{p,q}$ and $T_{p,q}$ to $S_{r,s}$ and $T_{r,s}$ if and only if $p + q + r + s$ is a special divisor on B , and λ exchanges $S_{p,q}$ and $T_{p,q}$*
- (5) *Two objects of $\mathcal{P}_5^{-1}(\mathcal{J}(B))$ are related by at most two tetragonal constructions.*

1.5. Theta Functions. Now, we shall describe one more major technical tool used in this paper. The *Riemann theta function* on $\mathbb{H}_g \times \mathbb{C}^g$ is given by

$$\theta(\Omega, z) = \sum_{n \in \mathbb{Z}^g} \exp[\pi i ({}^t n \Omega n + 2 {}^t n z)].$$

This function is periodic with respect to $\mathbb{Z}^g$ and is multiplied by an exponential factor with respect to $\Omega \mathbb{Z}^g$. Thus, the zero locus of $\theta(\Omega, z)$ is periodic for $\mathbb{Z}^g \oplus \Omega \mathbb{Z}^g$ and gives a divisor Θ on A , the *theta divisor*.

For $\epsilon, \delta \in \mathbb{Q}^g$, we define a *theta function with characteristics* to be

$$\theta \left[\begin{smallmatrix} \epsilon \\ \delta \end{smallmatrix} \right] (\Omega, z) = \exp[\pi i ({}^t \epsilon \Omega \epsilon + 2 {}^t \epsilon (z + \delta))] \theta(\Omega, z + \Omega \epsilon + \delta)$$

which is essentially the translate of $\theta(\Omega, z)$ by $\Omega\epsilon + \delta$. Evaluating at $z = 0$, these are Siegel modular forms of weight $\frac{1}{2}$ and level $(4, 8)$, so they are only well-defined on $\mathcal{A}_g(4, 8)$, not on $\mathcal{A}_g$ itself.

We will also need $\theta_2[\epsilon](\Omega, z) = \theta \begin{bmatrix} \epsilon \\ 0 \end{bmatrix} (2\Omega, 0)$ where $\epsilon \in (\frac{1}{2}\mathbb{Z}/\mathbb{Z})^g$, the second order theta functions, and $\theta \begin{bmatrix} \epsilon & 0 \\ 0 & 1/2 \end{bmatrix} (2\Omega, 0)$ where $\epsilon \in (\frac{1}{2}\mathbb{Z}/\mathbb{Z})^{g-1}$. These are modular forms of weight $(2, 4)$, and we can use them to define maps $\alpha_g: \mathcal{A}_g(2, 4) \rightarrow \mathbb{P}(U_g)$ and $\beta_g: \mathcal{RA}_g(2, 4) \rightarrow \mathbb{P}(U_{g-1})$ where U_g is the vector space of functions $(\mathbb{Z}/2\mathbb{Z})^g \rightarrow \mathbb{C}$ by setting $\alpha_g(\Omega)_\epsilon = \theta_2[\epsilon](\Omega, 0)$ and $\beta_g(\Omega)_\epsilon = \theta \begin{bmatrix} \epsilon & 0 \\ 0 & 1/2 \end{bmatrix} (2\Omega, 0)$.

We get maps on $\mathcal{A}_g$ and $\mathcal{RA}_g$ by noting that $G_g = \Gamma_g/\Gamma_g(2, 4)$ acts on $\mathcal{A}_g(2, 4)$, $\mathcal{RA}_g(2, 4)$ and $\mathbb{P}(U_g)$ in compatible ways, so that if $\mathbb{P}_g = \mathbb{P}(U_g)/G_g$, we have $\alpha_g: \mathcal{A}_g \rightarrow \mathbb{P}_g$ and $\beta_g: \mathcal{RA}_g \rightarrow \mathbb{P}_{g-1}$.

The primary purpose of this paper will be to understand the fibers of β_5 . For this, we will use

Theorem 1.19 (Theta symmetry [Don87b, Theorem 3.1]).

Let $C \in \mathcal{M}_{g+1}$ be a curve of genus $g+1$ and let $\{0, \mu_0, \mu_1, \mu_2\}$ be a rank 2 isotropic subgroup of $\mathcal{I}_{g+1}(C)_2$ (thus, $\mu_2 = \mu_0 + \mu_1$). For $i = 0, 1, 2$ we have a Prym variety $P_i = P(C, \mu_i) \in \mathcal{A}_g$ and on it a uniquely determined semiperiod ν_i , the image of μ_j , $j \neq i$ in P_i .

The point $\beta(P_i, \nu_i)$ is independent of $i = 0, 1, 2$.

We will also use the description of the fibers of β_4 that appears in the survey [Don88]

Proposition 1.20 ([Don88, Theorem 5.3]).

For $C \in \mathcal{M}_3$, the fiber $\beta_4^{-1}(\alpha_3(\mathcal{J}(C)))$ consists of two copies of the Kummer $K(\mathcal{J}(C))$, one contained in the interior of $\mathcal{RM}_4$ and the other the fiber over $\mathcal{J}(C) \in \mathcal{A}_3 = \partial\overline{\mathcal{A}}_4^s$ in the projection $\overline{\mathcal{A}}_4^t \rightarrow \overline{\mathcal{A}}_4^s$.

1.6. The Schottky Loci. We begin with the theorem that motivates this study:

Theorem 1.21 (Schottky-Jung Identities [Sch88, SJ09, RF74]).

The following diagram commutes:

$$\begin{array}{ccc}
 & \mathcal{RM}_g & \\
 \mathcal{P}_g \swarrow & & \searrow \mathcal{RJ}_g \\
 \mathcal{A}_{g-1} & & \mathcal{RA}_g \\
 \alpha_{g-1} \searrow & & \swarrow \beta_g \\
 & \mathbb{P}_g &
 \end{array}$$

Schottky and Jung noticed these relations between various theta functions and suggested that they can be used to describe Jacobians. More precisely, we define

$$\begin{aligned}
 \mathcal{RS}_g &= \beta_g^{-1}(\text{im } \alpha_{g-1}) \\
 \mathcal{S}_g^{\text{big}} &= \{A | \exists \mu, (A, \mu) \in \mathcal{RS}_g\} \\
 \mathcal{S}_g^{\text{small}} &= \{A | \forall \mu, (A, \mu) \in \mathcal{RS}_g\}
 \end{aligned}$$

Schottky and Jung conjectured that $\mathcal{S}_g^{\text{big}} = \mathcal{J}_g$, the closure of the image of the Torelli map. This is known not to be true, and the key result for that is Theta symmetry (Theorem 1.19)

Let $Q \subset \mathbb{P}^2$ be a plane quintic curve and let μ, ν be two points of order two on $\mathcal{J}(Q)$ such that μ is odd and ν is even. Then $(\mathcal{P}(Q, \mu), \bar{\nu}) \in \mathcal{RS}_5$ if and only if $(\mathcal{P}(Q, \nu), \bar{\mu}) \in \mathcal{RS}_5$. However, as mentioned above and proved in [DS81], $\mathcal{P}(Q, \mu)$ is the intermediate Jacobian of a cubic threefold. Additionally, $\mathcal{P}(C, \nu)$ is known to be a Jacobian of a curve (see the discussion in section 5 of [Don87b]). Thus, $\mathcal{C} \subset \mathcal{S}_5^{\text{big}}$, and so Donagi introduced $\mathcal{S}_g^{\text{small}}$ to correct this, as only $\mathcal{RC}^0$ appears in $\mathcal{RS}_5$, not $\mathcal{RC}^1$, the locus of intermediate Jacobians of cubic threefolds with an odd point of order two.

There are two additional components that are not difficult to show lie in $\mathcal{RS}_5$. There is the locus $\mathcal{RA}_1 \times \mathcal{A}_4$, and in fact:

Proposition 1.22 ([Don87a, 3.3.4]).

The Schottky locus $\mathcal{RS}_g$ contains $\mathcal{A}_{g'} \times \mathcal{RA}_{g''}$ for all $g' + g'' = g$ (Note that if $g = 0$, then $\mathcal{RS}_0 = \emptyset$).

We will see below that no other product loci can be components of $\mathcal{RS}_5$.

There is also one boundary component, described by

Proposition 1.23 ([Don87a, Theorem 3.3.1]).

In the Satake compactification, we have

$$\partial \overline{\mathcal{RS}}_g^s = \partial^I \overline{\mathcal{RA}}_g^s \cup \partial^{III} \overline{\mathcal{RA}}_g^s \cup i_{II}(\mathcal{RS}_{g-1})$$

where i_{II} is the inclusion of $\mathcal{RA}_{g-1}$ as $\partial^{II} \overline{\mathcal{RA}}_g^s$.

In [Don88] this result is strengthened to show that $\partial \overline{\mathcal{RA}}_g^t$ does not contain ∂^{III} or $i_{II}(\mathcal{RS}_{g-1})$ as components, but only contains the points that are limits of other components.


2. LINE CONFIGURATIONS

In this section, we will study a class of configurations that occur whenever there is a triality on a space, such as the tetragonal construction or theta symmetry. Because we are working with trialities, everything in this section will be done over $\mathbb{F}_2$, but the majority will work for $\mathbb{F}_q$.

Definition 2.1 (Line Configuration).

A line configuration V over $\mathbb{F}_2$ is a set P_V , called the points of V , along with a set L_V , called the lines of V , such that for all $\ell \in L_V$, we have $\ell \subset P_V$, $|\ell| = 3$ and for all $\ell, \ell' \in L_V$, $|\ell \cap \ell'| \geq 2$ implies that $\ell = \ell'$.

We can construct a large class of examples, which we will call *algebraic line configurations*. These are defined by starting with $V \subset \mathbb{P}^n$ over $\mathbb{F}_2$ a projective variety. Then we set P_V to be the $\mathbb{F}_2$ -points of V and L_V to be the projective lines over $\mathbb{F}_2$ contained in V . The first interesting example is $V = \mathbb{P}^2$, the Fano plane:



we define a morphism of configurations $f: V \rightarrow W$ to be an injection $f: P_V \rightarrow P_W$ such that for each line $\{p_0, p_1, p_\infty\} \in L_V$, the image $\{f(p_0), f(p_1), f(p_\infty)\}$ is a line in L_W . A *subconfiguration* is then the image of a morphism. We say that two points are collinear if they lie on a subconfiguration isomorphic to $\mathbb{P}^1$ and that two lines are coplanar if they lie on a subconfiguration isomorphic to $\mathbb{P}^2$.

Definition 2.2 (V-Configuration).

Let V be a line configuration. Then another line configuration W is a *V-configuration* if for each $p \in P_W$, there exists a bijection $\phi_p: \{\ell \in L_W | p \in \ell\} \rightarrow P_V$ such that ℓ, ℓ' are coplanar if and only if $\phi_p(\ell)$ and $\phi_p(\ell')$ are collinear.

Example 2.3.

- (1) $\mathbb{P}^n$ is a $\mathbb{P}^{n-1}$ -configuration.
- (2) If V is a collection of n points with $L_V = \emptyset$, then $(\mathbb{P}^1)^n$ is a V-configuration.
- (3) For some V , we can construct examples of multiple fundamentally distinct V-configurations. For instance, if V consists of five points and no lines, the previous example says that $(\mathbb{P}^1)^5$ is a V-configuration. However, if S is a smooth cubic surface, then we can define a configuration W with $P_W = \{\ell \subset S | \ell \text{ is a line}\}$ and with L_W the set of triples of coplanar lines. It is classical that given a line on a cubic surface, it is contained in exactly five coplanar triples, no set of which are configured as a Fano plane.

Now, to help us to describe the properties of line configurations, we use the *incidence graph*. This is the graph Γ_V whose vertices are the points of P_V and two vertices $p, q \in P_V$ are connected by an edge if and only if they are collinear. We fix a metric on Γ_V such that each edge has length 1, and give properties to V from the properties of the metrized graph Γ_V , for instance, we can speak of connected configurations or the diameter of a configuration.

Given Γ_V , we can define numerical invariants of a configuration. For each $p \in P_V$, we define $V_i(p) = \{q \in V | d(p, q) = i\}$, the points that are first reached after passing along i lines from p . For points $p \in P_V$, and $q \in V_i(p)$, we define $V_{i,j}(p, q) = V_j(p) \cap V_1(q)$, the points distance j from p which are adjacent to q . Note that $V_{i,j}(p, q)$ is empty unless j is $i-1$, i or $i+1$. We denote the cardinalities of these sets by $v_i(p)$ and $v_{i,j}(p, q)$ (by convention, we set these numbers to be zero if i or j is negative), and call a line configuration *symmetric* if the $v_i(p)$ and $v_{i,j}(p, q)$ don't depend on p and q , in which case we will denote $v_i = v_i(p)$ and $v_{i,j} = v_{i,j}(p, q)$.

Proposition 2.4.

Let V be a connected symmetric line configuration. Then

- (1) For all i , $v_1 = v_{i,i-1} + v_{i,i} + v_{i,i+1}$
- (2) For all i , $v_1 v_i = v_{i-1,i} v_{i-1} + v_{i,i} v_i + v_{i+1,i} v_{i+1}$
- (3) We have $v_0 = 1$, $v_{0,0} = 0$, $v_{0,1} = v_1$ and $v_{1,0} = 1$.

Let W be a connected symmetric V-configuration. Then

- (4) $w_{2,2} \geq w_{2,1}$
- (5) $w_1 = 2|P_V|$
- (6) $w_{1,1} = 2v_1 + 1$
- (7) If, additionally, $v_3 = w_3 = 0$, then either $w_2 = |P_V|$ or $w_2 = 4v_2$.

For convenience, throughout the proof we will set $\text{Lines}_W(p) = \{\ell \in L_W | p \in \ell\}$, the set of lines through p in W .

Proof. Let V be a connected symmetric line configuration and W a connected symmetric V -configuration.

- (1) Fix $p \in V$, $i \in \mathbb{N}$, $q \in V_i(p)$. Then

$$\begin{aligned} & V_{i,i-1}(p, q) \cup V_{i,i}(p, q) \cup V_{i,i+1}(p, q) \\ &= (V_{i-1}(p) \cap V_1(q)) \cup (V_i(p) \cap V_1(q)) \cup (V_{i+1}(p) \cap V_1(q)) \\ &= (V_{i-1}(p) \cup V_i(p) \cup V_{i+1}(p)) \cap V_1(q) \\ &= V_1(q). \end{aligned}$$

- (2) Fix $p \in V$, $i \in \mathbb{N}$. Let $X = \{(a, b) | a \in V_i(p), b \in V_1(a)\}$. Then $|X| = v_i v_1$. But also,

$$\begin{aligned} X &= \{(a, b) | a \in V_i(p) \cap V_1(b), b \in \cup_j V_j(p)\} \\ &= \cup_j \{(a, b) | a \in V_{j,i}(p, b), b \in V_j(p)\} \end{aligned}$$

and so $|X| = \sum_j v_{j,i} v_j$.

- (3) These all follow directly from the definitions.
- (4) Fix $q \in W_2(p)$. We prove that no line containing q contains two points of $W_1(p)$. As each line consists of 3 points, this implies that $w_{2,2} \geq w_{2,1}$. Let $a, b \in W_1(p)$ and assume that there is a line $\ell \in L_W$ such that $a, b, q \in \ell$. As $a, b \in W_1(p)$, there exist lines m_1, m_2 through p containing a, b respectively. But then, m_1, m_2 must be coplanar, and so there is a line m containing p, q , so $q \in V_1(p) \cap W_2(p) = \emptyset$, a contradiction.
- (5) For each line $\ell \in \text{Lines}_W(p)$, fix a bijection $\ell \rightarrow \mathbb{P}^1(\mathbb{F}_2)$ such that p is mapped to ∞ . Then $\cup_{\ell \in \text{Lines}_W(p)} \ell \setminus \{p\} = \coprod_{\ell \in \text{Lines}_W(p)} \mathbb{A}^1(\mathbb{F}_2)$, and this has cardinality twice the number of lines, $2|P_V|$.
- (6) Fix $p \in P_W$, ℓ a line through p , $p' \in \ell$ distinct from p . Through p' , there are $|P_V|$ lines. One is ℓ , v_1 of them are coplanar with ℓ , and the rest are not. Each coplanar line consists of n points in $V_1(p)$ that are not p' , but there are also $n-1$ points of ℓ in $V_1(p)$ other than p' , and so $w_{1,1} = 2v_1 + 1$.
- (7) Fix $p \in W$. Let $\tilde{X}$ be the set of triples (ℓ, m, q) in $\text{Lines}_W(p) \times L_W \times W_2$ such that $p \in \ell$, $\ell \cap m \neq \emptyset$ and $q \in m$. There is a natural map $\tilde{X} \rightarrow W_2$ which is surjective. Then

$$\begin{aligned} |\tilde{X}| &= w_2 \cdot |\text{fiber}| \\ &= w_2 \cdot |\{\text{paths to } q \in W_2 \text{ from } p\}| \\ &= w_2 \cdot |\{\text{points of } W_1(p) \text{ connected to } p\}| \\ &= w_2 \cdot |W_1(p) \cap W_1(q)| \text{ for } q \in W_2(p) \\ &= w_2 w_{2,1} \end{aligned}$$

Now, we also have a map $\tilde{X} \rightarrow \text{Lines}_W(p) \times W_2(p) = A \coprod B$, where the fiber over A has cardinality 1 and over B has cardinality 0. These are the only possibilities, because if there were two, then we get a plane and $q \in W_1(p)$. So $|A| = |\tilde{X}| = w_2 w_{2,1}$, and $|A| + |B| = |\text{Lines}_W(p) \times W_2(p)| = |P_V| w_2$, so $|B| = w_2(|P_V| - w_{2,1})$.

But, as W is symmetric, we can see that $|V| = |P_V| \alpha$, where α is the number of lines in $\text{Lines}_W(p)$ that don't have a line connecting them to q . Then $\alpha = |P_V| - |\{\text{lines in } \text{Lines}_W(p) \text{ connected to } q\}|$, which is $\alpha = |V| - w_{2,1}$, so $|B| = |P_V|(|P_V| - w_{2,1})$.

So, $w_2(|P_V| - w_{2,1}) = |P_V|(|P_V| - w_{2,1})$, implying that either $w_2 = |P_V|$ or else $w_{2,1} = |P_V|$. In the latter case, the previous parts of this proposition along with the hypothesis that $\text{diam}(W) \leq 2$, implies that $w_2 = 4v_2$. $\square$

We can now apply these numerics to another family. Let the zero locus of $x_1^2 + x_2^2 + x_1x_2 + x_3x_4 + \dots + x_{2n-1}x_{2n}$ in $\mathbb{P}^{2n-1}(\mathbb{F}_2)$ be denoted by Q_{2n}^- . This is the smooth quadric of Witt defect 1 in $\mathbb{P}^{2n-1}$ over $\mathbb{F}_2$. We have actually already seen it in an above example: if $n = 2$, then the zero locus is five points, no three collinear, and if $n = 3$, we get 27 points, each of which lies on 5 lines (and thus is adjacent to 10 points), which gives the configuration of lines on a smooth cubic surface. The latter fact also appears as an isomorphism $WE_6 \cong O_6^-(\mathbb{F}_2)$, the orthogonal group preserving the form $x_1^2 + x_2^2 + x_1x_2 + x_3x_4 + x_5x_6$.

For all n , in fact, we can see that Q_{2n+2}^- is a Q_{2n}^- -configuration, by noting that they are homogeneous varieties, and so we only need to look at the point $[0 : \dots : 0 : 1]$, where everything is easy to compute. Additionally, for $n \geq 3$ these are all symmetric and have diameter two, so all of the numerical conditions above apply.

Lemma 2.5.

Let $n \geq 2$. Then the only connected, symmetric Q_{2n}^- -configuration of diameter 2 is Q_{2n+2}^- .

Proof. We set $V = Q_{2n}^-$ for some fixed $n \geq 2$ and W a connected, symmetric V -configuration.

As we are assuming diameter 2, we have to determine w_i and $w_{i,j}$ for $0 \leq i, j \leq 2$. Proposition 2.4(3) determines $w_0, w_{0,0}, w_{1,0}$ and that $w_{0,1} = w_1$. Then Proposition 2.4(5) says that $w_1 = 2|Q_{2n}^-| = 2(2^{n-1}(2^n - 1) - 1) = 2^n(2^n - 1) - 2 = 2^{2n} - 2^n - 2$.

This leaves $w_{1,1}, w_{2,1}, w_{1,2}, w_{2,2}$ and w_2 . Proposition 2.4(6) gives $w_{1,1} = 2(2^{2n-2} - 2^{n-1} - 2) = 2^{2n-1} - 2^n - 4$. Then Proposition 2.4(1) says that $2^{2n} - 2^n - 2 = 1 + w_{1,1} + w_{1,2}$ and $2^{2n} - 2^n - 2 = w_{2,1} + w_{2,2}$, which we solve for $w_{1,2} = 2^{2n} - 2^n - 2 - 1 - (2^{2n-1} - 2^n - 4) = 2^{2n} - 2^{2n-1} + 1$.

Substitution of 2.4(1) into 2.4(2) and cancellation gives us the relation $w_i w_{j,i} = w_j w_{i,j}$. And thus, $w_1 w_{2,1} = w_2 w_{1,2}$. So $(2^{2n-1} - 2^n - 4)w_{2,1} = w_2(2^{2n} - 2^n - 2 - 1 - (2^{2n-1} - 2^n - 4))$. Using 2.4(7), we know that $w_2 = (2^{n-1}(2^n - 1) - 1)$ or $4((2^{n-1}(2^n - 1) - 1) - 1 - 2((2^{n-2}(2^{n-1} - 1) - 1))) = 2^{2n}$. The former case leads to $w_{2,2} < w_{2,1}$ contradicting 2.4(4), and so we must have the latter case. And finally, this determines $w_{2,1}$ using $w_1 w_{2,1} = w_2 w_{1,2}$ and $w_{2,2}$ from $w_1 = w_{2,1} + w_{2,2}$.

In particular, $|W| = |Q_{2n+2}^-| = 2^n(2^{n+1} - 1) - 1$.

We now prove that these $2^n(2^{n+1} - 1) - 1$ points must be connected by lines in a unique way to satisfy the conditions of the lemma. We begin with a point of the configuration $w \in W$. As $w_{2,1} = w_{2,2} = 2^{n-1}(2^n - 1) - 1 = |Q_{2n}^-|$, each line not containing w must contain one point in $W_1(w)$ and two points of $W_2(w)$. Thus, a point in $W_2(w)$ is given by a point (not w) on each line containing w .

The same holds for any point $w' \in W_1(w)$, and there must be a function $W_1(w) \rightarrow W_1(w')$ that preserves incidence and takes the set of choices at w to those at w' . This will dictate the data of which half (we note that $w_{1,2} = \frac{1}{2}w_2$, and so each point of $W_1(w)$ is colinear with half of the points of $W_2(w)$) of the points of $W_2(w)$ are connected to w' . As this works for each $w' \in W_1(w)$, there is only one

such choice that will work globally, up to automorphisms of Q_{2n}^- and involutions switching non- w points on a line. $\square$

So to identify a Q_{2n}^- -configuration, we must merely check symmetry and diameter.

Proposition 2.6.

The general fiber of $\beta_5: \mathcal{RA}_5 \rightarrow \mathbb{P}^{15}/G_4$ is a Q_6^- -configuration.

Proof. Let $(A, \mu) \in \mathcal{RA}_5$ be a general point. Then A is the Prym variety of 27 distinct curves $(X_i, \nu_i) \in \mathcal{RM}_6$, and μ lifts to two distinct points of order two on $\mathcal{J}(X_i)$, μ_i^0, μ_i^1 . Then theta symmetry implies that, again generically, $(\mathcal{P}(X_i, \mu_i^j), \bar{\nu}_i)$ are distinct points in the fiber of β_5 . Thus, (A, μ) lies on 27 triples $(A, \mu), (\mathcal{P}(X_i, \mu_i^0), \bar{\nu}_i), (\mathcal{P}(X_i, \mu_i^1), \bar{\nu}_i)$, which are lines over $\mathbb{F}_2$, form the configuration Q_6^- , by Theorem 1.10, and so the fiber is a Q_6^- -configuration. $\square$

Lemma 2.7.

The connected components of the Q_6^- -configuration on the fibers of β_5 has diameter 2.

Proof. If we blow up $\mathcal{RP}_6: \mathcal{R}^2\mathcal{M}_6 \rightarrow \mathcal{RA}_5$ to a finite map $\widetilde{\mathcal{RP}}_6$, then Proposition 1.12 says that the fiber over $(X, \mu) \in \mathcal{RC}^0$, with the normal data suppressed for ease of notation, consists of 54 plane quintics with an odd and an even point of order two marked. Theta symmetry then says that the Pryms with respect to the even points with the images of the odd points marked are in the same fiber of β_5 , and all of these are the Jacobians of curves. By Proposition 1.13, the fiber of a Jacobian consists of quintic curves, whose Pryms are cubic threefolds and Jacobians, trigonal curves, whose Pryms are Jacobians, and singular curves, whose Pryms are Jacobians and degenerate abelian varieties in $\partial^I \overline{\mathcal{RA}}_5^t$.

If we start with a degenerate abelian variety, then we must only have singular curves lying over it. Proposition 1.4 tells us that these are either in ∂^{II} or $\partial_{i,6-i}$. But the latter component only gives products, and so generically all 27 preimages under $\mathcal{P}_6$ are in ∂^{II} . Thus, any points of $\overline{\mathcal{RA}}_5^t$ related to a point of $\partial^I \overline{\mathcal{RA}}_5^t$ is either in $\partial^I \overline{\mathcal{RA}}_5^t$ or is a Jacobian, and so no new types of points occur after the second iteration of theta symmetry.

Finally, we show that the objects obtained previously don't constitute anything new. First, the Jacobians obtained from the chosen Jacobian in the second step. These Jacobians all must have already been obtained from the cubic threefold, because each Jacobian is the Prym of a unique quintic curve, and will thus determine a cubic threefold. However, generically no two cubic threefolds can be related in this manner, because the incidence is contained in the fibers of β_5 , which induces a birational isomorphism between $\mathcal{RC}^0$ and $\alpha(\mathcal{A}_4)$, and so, generically, must be injective. And last, we have the objects obtained from a given degenerate abelian variety. We have twenty-seven nodal curves, and each occurs as a Wirtinger and as a ∂^{II} double cover, which we then take the Pryms of. The Wirtingers must be Jacobians we've already seen, by the above, and so the ∂^{II} double covers would be obtained by applying theta symmetry on those Jacobians first. Thus, the Q_6^- -configuration must have diameter 2. $\square$

And so, by Lemma 2.5, the map β_5 must have degree a multiple of 119.

3. DEGENERATIONS OF ABELIAN VARIETIES

In this chapter, we will show that the Q_6^- -configuration on the fibers of β is connected. This amounts to computing the degree of β , which we will do by studying certain degenerations. We will spend this chapter studying the structure of $\partial^{II}\overline{\mathcal{RA}}_5^t$, the degenerations with vanishing cycle orthogonal but not identical to the marked semiperiod.

Lemma 3.1 ([Don87a, Corollary 3.2.3 and Lemma 3.3.6]).

The extension of $\beta_g: \overline{\mathcal{RA}}_g^t \rightarrow \mathbb{P}(U_{g-1})/G_{g-1}$ to $\partial^{II}\overline{\mathcal{RA}}_g^t$ is β_{g-1} and the diagram:

$$\begin{array}{ccc} \mathcal{RA}_{g-1} & \xrightarrow{i_{II}} & \overline{\mathcal{RA}}_g^t \\ \beta_{g-1} \downarrow & & \downarrow \beta_g \\ \mathbb{P}(U_{g-2})/G_{g-2} & \longrightarrow & \mathbb{P}(U_{g-1})/G_{g-1} \end{array}$$

is Cartesian.

This result tells us that if we start with a point in $\partial^{II}\overline{\mathcal{RA}}_g^t$ then the whole fiber over its image is in $\partial^{II}\overline{\mathcal{RA}}_g^t$, and so we focus our attention there:

Proposition 3.2.

Any two points in a fiber of β_5 inside $\partial^{II}\overline{\mathcal{RA}}_5^t$ are related by a sequence of theta symmetries.

Proof. By Lemma 3.1, we are working with the map $\beta_4: \mathcal{RA}_4 \rightarrow \mathbb{P}^7/G_3$. Now, fix a curve $C \in \mathcal{M}_3$. Then the fiber of β_4 over $\alpha_3(\mathcal{J}(C))$ is $\text{Bl}_0 K(\mathcal{J}(C)) \cup K(\mathcal{J}(C))$ by Proposition 1.20, with the first component consisting of genus 4 Jacobians and the second of degenerate abelian varieties in $\partial^I\overline{\mathcal{RA}}_4^t$. Over $(B, \mu) \in \mathcal{RJ}_4$, the fiber of $\mathcal{RP}_5$ is a double cover of $\text{Sym}^2 B \cup \text{Sym}^2 B$, with the first component consisting of trigonal curves and the second consisting of Wirtinger curves, as in Proposition 1.18. Thus, for each $(B, \tilde{B})$, we get a map from the double cover of $\text{Sym}^2 B, \text{Sym}^2 B \rightarrow \text{Bl}_0 K(\mathcal{J}(C))$, because theta symmetry takes trigonal curves to trigonal curves and Wirtinger curves to Wirtinger curves. Each of these maps has two dimensional image, and they're all nonisomorphic, and thus distinct, so the dimension of the union is at least three, so surjectivity follows. $\square$

Now that we have a locus where we know that theta symmetry spans the fibers, we need to identify theta symmetry over that locus:

Proposition 1.15 tells us that the fiber in $\partial^{II}\overline{\mathcal{RA}}_5^t$ has the Q_6^- -configuration structure we expect, and more so, the fact that theta symmetry spans the fibers implies connectivity, so the fiber is Q_8^- . It remains, though, to check that β_5 is generically unramified on $\partial^{II}\overline{\mathcal{RA}}_5^t$.

We have a tower of maps $\overline{\mathcal{R}^2\mathcal{M}}_6 \xrightarrow{\mathcal{RP}_6} \overline{\mathcal{RA}}_5^t \xrightarrow{\beta_5} \mathbb{P}^{15}/G_4$ and as each map is generically finite, if we blow up each locus with positive dimensional fibers, and the blow up again so that the maps are all well defined, the maps will be finite. We denote the blowups and blown up maps by $\widetilde{\mathcal{R}^2\mathcal{M}}_6 \rightarrow \widetilde{\mathcal{RA}}_5 \rightarrow \widetilde{\mathbb{P}}$. Set $X = \widetilde{\mathcal{RA}}_5 \times_{\mathbb{P}} \widetilde{\mathcal{RA}}_5$. For each point $(x, y) \in X$, we can associate the distance in $\Gamma_{Q_8^-}$, or ∞ if they are not connected. This gives a decomposition $X = I_1 \cup I_{54} \cup I_{64} \cup I$,

where I_1 is the diagonal, I_{54} are theta related pairs, I_{64} the pairs theta related to a common point and I the pairs in the same fiber but not theta related. Our goal is to show that $I = \emptyset$.

We will focus on studying I_{54} . It fits into a diagram

$$\begin{array}{ccc}
 I_{54} & \xrightarrow{\tilde{\beta}} & \widetilde{\mathcal{RA}}_5 \\
 \downarrow 2:1 & \nearrow \tilde{\mathcal{P}} & \\
 I_{27} & &
 \end{array}$$

where I_{27} is the pullback of the Prym map to $\widetilde{\mathcal{RA}}_5$ from $\mathcal{A}_5$, and $I_{54} \rightarrow I_{27}$ is the natural double cover. Differentiating these maps, we can see that $\text{Ram } \tilde{\beta} = \text{Ram } \tilde{\mathcal{P}}$.

Lemma 3.3.

Let $f: X \rightarrow Y$ be a finite morphism of smooth varieties and $\bar{f}: X \times_Y X \rightarrow X$ be the fiber product of f with itself. Then if p is a ramification point of f and q is a nonramification point in the same fiber, (p, q) is a ramification point of $\bar{f}$.

Proof. Set $b = f(q) = f(p)$. As p is a ramification point and q is not, we have df_q an isomorphism and df_p not. Locally, this means that we have an isomorphism of X and Y near q but not near p , and so the map $d\bar{f}_{(p,q)}: T_{p,q}(X \times_Y X) \rightarrow T_p X$ is not an isomorphism. $\square$

Lemma 3.4.

Let $p \in \partial^{II} \widetilde{\mathcal{RA}}_5$. Then if β_5 is ramified at p , I_{54} is ramified over p .

Proof. Let $p \in \partial^{II} \widetilde{\mathcal{RA}}_5$ be a ramification point of $\tilde{\beta}$. Then either there exists a q such that $(p, q) \in I_{54}$ is unramified, or else not. If there is such a q , then this follows from Lemma 3.3.

If not, then we must have a fiber of the Prym map which is totally ramified. This would correspond to a cubic surface such that every line has multiplicity at least two, but which is reduced and irreducible. This cannot happen, as if there is a line of multiplicity two, then the residual intersection with the tangent plane along that line must be a line of multiplicity one, so every line would have multiplicity three, which does not occur. $\square$

More useful than the lemma is the contrapositive: that if I_{54} is unramified over p , then β_5 is unramified at p .

Proposition 3.5.

The map β_5 is generically unramified on $\partial^{II} \widetilde{\mathcal{RA}}_5^t$.

Proof. Lemma 3.4 tells us that a point is a ramification point for β_5 only if it is a branch point of $\tilde{\beta}$, which is the same as being a branch point over $\tilde{\mathcal{P}}$. But Izadi's Theorem 1.15 tells us that the Prym map is generically unramified over $\partial^{II} \widetilde{\mathcal{RA}}_5^t$, and so $I_{54} \rightarrow \widetilde{\mathcal{RA}}_5$ is generically unramified over $\partial^{II} \widetilde{\mathcal{RA}}_5^t$, and so, β_5 is as well. $\square$

Thus, we have the main result of this section:

Theorem 3.6.

the fibers of $\beta_5: \overline{\mathcal{RA}}_5^t \rightarrow \mathbb{P}^{15}/G_4$ are exactly the orbits of theta symmetry.

4. CONTRACTED LOCI

Now, we've identified the fibers of β_5 with the orbits of theta symmetry, which gives us the degree of this generically finite map. What remains is to understand the locus where β_5 has infinite fibers. As each abelian variety has finitely many points of order two, and as α_g is finite and generically injective for all g [SM94, Proposition 1], any infinite fiber must arise from an infinite fiber of the Prym map.

By Proposition 1.11 we know that given $(C, \mu) \in \mathcal{RM}_6$, we obtain the rest of the fiber containing (C, μ) by iterating the tetragonal construction, so long as we allow points to be related via the boundary. There can be no infinite chains of curves with finitely many g_4^1 's, because such a fiber would be countable, and an infinite fiber must be uncountable. Thus, infinite fibers will correspond to curves with infinitely many g_4^1 's and to singular curves.

Theorem 4.1 ([ACGH85, Theorem IV.5.2], [Mum74, Theorem in Appendix]).

Let C be a smooth non-hyperelliptic curve of genus $g \geq 4$. Let $d, r \in \mathbb{Z}$ with $0 < 2r \leq d$ and $2 \leq d \leq g - 2$. Then, if there exists a $(d - 2r - 1)$ -dimensional family of g_d^r 's on C , we must have that C is trigonal, bielliptic or a plane quintic.

Fixing $g = 6$, $d = 4$ and $r = 1$, the hypotheses on Theorem 4.1 are satisfied, and so a curve of genus 6 that has a positive dimensional family of g_4^1 's must be hyperelliptic, trigonal, bielliptic, a plane quintic, or singular.

Proposition 1.9 says that Pryms of hyperelliptic curves are products. Theorem 1.8 tells us that trigonal Pryms are Jacobians. Propositions 1.12 and 1.13 tell us that Pryms of quintics are Jacobians or cubic threefolds and Proposition 1.4 says that the Prym varieties of singular curves are contained in $\mathcal{J}_5$, $\partial \overline{\mathcal{A}}_5^t$, the Beauville Pryms or the product loci.

But we actually only care about contracted loci over $\alpha_4(\mathcal{A}_4)$, which may be irreducible components of $\mathcal{RS}_5$. As $\dim(\alpha_4(\mathcal{A}_4)) = 10$, any irreducible component of $\mathcal{RS}_5$ must have dimension at least 10. Also, as we will see later, $\overline{\mathcal{RJ}}_5$, $\overline{\mathcal{RC}}^0$, and $\partial^I \overline{\mathcal{RA}}_5^t$ are the only components of $\mathcal{RS}_5$ with positive local degree, thus, any other component is actually a contracted locus.

Proposition 4.2.

The only component of $\beta_5^{-1}(\alpha_4(\mathcal{A}_4))$ that is blown down by β is $\mathcal{RA}_1 \times \mathcal{A}_4$.

Proof. By the above discussion, the only loci with positive dimensional fibers are $\mathcal{RJ}_5$, the bielliptic loci, $\mathcal{RC}^0$, $\mathcal{RC}^1$, $\partial^I \overline{\mathcal{RA}}_5^t$, $\partial^{II} \overline{\mathcal{RA}}_5^t$, $\partial^{III} \overline{\mathcal{RA}}_5^t$, and $\mathcal{RP}_6(\partial^{III} \overline{\mathcal{R}^2 \mathcal{M}}_6)$.

The loci $\mathcal{RJ}_5$, $\mathcal{RC}^0$ and $\partial^I \overline{\mathcal{RA}}_5^t$ actually have positive local degree (see next chapter) and so are not blown down.

The bielliptic loci are ruled out by 1.6.

The locus $\mathcal{RC}^1$ is not in $\mathcal{RS}_5$, as any point of $\mathcal{RC}^1$ will only be theta related to other points of $\mathcal{RC}^1$. Specifically, a curve over $(X, \mu) \in \mathcal{RC}^1$ must be a plane quintic Q and an odd point of order two ν with $P(Q, \nu) = X$, but also the lifts of μ must be odd, so all three points in any instance of theta symmetry give points of $\mathcal{RC}^1$.

From 1.23, it is shown that $\partial^{III}\overline{\mathcal{RA}}_5^t$, the Beauville Pryms, $\mathcal{RA}_4 \times \mathcal{A}_1$ and $\mathcal{RA}_4 \times \mathcal{RA}_1$ are not in the Schottky locus, and all other product loci other than $\mathcal{RA}_1 \times \mathcal{A}_4$ are of too small dimension. However, $\mathcal{RA}_1 \times \mathcal{A}_4$ is contracted by β_5 .

The only remaining component is $\partial^{II}\overline{\mathcal{RA}}_5^t$, but we know this locus is mapped to $\mathbb{P}^7/G_3$, and so any points of it in $\mathcal{RS}_5$ must be the limits of points of other components. $\square$

5. SCHOTTKY-JUNG LOCUS

Finally, we prove our main theorem:

Theorem 5.1.

In $\overline{\mathcal{RA}}_5^t$, we have $\overline{\mathcal{RS}}_5 = \overline{\mathcal{RJ}}_5 \cup \overline{\mathcal{RC}}^0 \cup \partial^I\overline{\mathcal{RA}}_5^t \cup \mathcal{A}_4 \times \mathcal{RA}_1$.

Proof. As the general fiber of β is a connected Q_6^- -configuration, we have that $\deg \beta = 119$. Then Proposition 4.2 tells us that the only blown down component of $\mathcal{RS}_5$ is $\mathcal{A}_4 \times \mathcal{RA}_1$, so it remains to identify the components with positive local degree.

It is known Proposition 1.16 that β_5 restricts to $\chi^{-1} : \mathcal{RC}^0 \dashrightarrow \mathcal{A}_4$, and thus has local degree one.

The Schottky-Jung relations imply that the restriction to $\overline{\mathcal{RJ}}_5$ is birational to the Prym map $\mathcal{RM}_5 \rightarrow \mathcal{A}_4$. Thus, over $A \in \mathcal{A}_4$, we get a double cover of the Fano surface of a cubic threefold. However, when we blow up to obtain a finite map, we get a double cover of the 27 lines on a cubic surface, and so we have local degree 54.

In [GG86], van Geemen and van der Geer compute the local degree on $\partial^I\overline{\mathcal{RA}}_5^t$ as follows: fix $X \in \mathcal{A}_4$. Then the part of the fiber of β_5 is $K(X)$ and by blowing up, they showed that the degree of the map is the same as that of the map $K(X) \rightarrow \mathbb{P}^4$ given by the linear system $\Gamma_{00} = \{s \in \Gamma(X, 2\Theta) \mid \text{mult}_0 s \geq 4\}$. We blow X up at 0, call the exceptional divisor E , and then this linear system is precisely $2\Theta - 4E$ on the abelian variety. Thus, the local degree is $\frac{1}{2}(2\Theta - 4E)^4 = 64$.

This gives us degree $1 + 54 + 64 = 119$, and so there are no other components with positive local degree. $\square$

And so, as noted in [Don87b], this implies

Corollary 5.2.

$$\mathcal{S}_5^{\text{small}} = \mathcal{J}_5$$

Proof. The points of $\mathcal{S}_5^{\text{small}}$ are just the abelian varieties A such that $(A, \mu) \in \mathcal{RS}_5$ for all nonzero points of order two $\mu \in A$. Among the components above, Jacobians have all nonzero points of order two, but the intermediate Jacobians of cubic threefolds only have even points of order two, the boundary component has only the vanishing cycle, and the product locus only has pullbacks of the points of order two on the elliptic curve. $\square$

REFERENCES

- [ACGH85] Enrico Arbarello, M. Cornalba, Phillip A. Griffiths, and Joseph Harris. *Geometry of algebraic curves. Vol. I*, volume 267 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1985.
- [AM67] Aldo Andreotti and A. L. Mayer. On period relations for abelian integrals on algebraic curves. *Ann. Scuola Norm. Sup. Pisa (3)*, 21:189–238, 1967.

- [AMRT10] Avner Ash, David Mumford, Michael Rapoport, and Yung-Sheng Tai. *Smooth compactifications of locally symmetric varieties*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010. With the collaboration of Peter Scholze.
- [Bea77a] Arnaud Beauville. Prym varieties and the Schottky problem. *Invent. Math.*, 41(2):149–196, 1977.
- [Bea77b] Arnaud Beauville. Variétés de Prym et jacobiniennes intermédiaires. *Ann. Sci. École Norm. Sup. (4)*, 10(3):309–391, 1977.
- [Bea89] Arnaud Beauville. Prym varieties: a survey. In *Theta functions—Bowdoin 1987, Part 1 (Brunswick, ME, 1987)*, volume 49 of *Proc. Sympos. Pure Math.*, pages 607–620. Amer. Math. Soc., Providence, RI, 1989.
- [CG00] Ciro Ciliberto and Gerard van der Geer. The moduli space of abelian varieties and the singularities of the theta divisor. In *Surveys in differential geometry*, *Surv. Differ. Geom.*, VII, pages 61–81. Int. Press, Somerville, MA, 2000.
- [Cle83] C. Herbert Clemens. Double solids. *Adv. in Math.*, 47(2):107–230, 1983.
- [CM08] Sebastian Casalaina-Martin. Singularities of theta divisors in algebraic geometry. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 25–43. Amer. Math. Soc., Providence, RI, 2008.
- [Deb90] Olivier Debarre. Variétés de Prym et ensembles d’Andreotti et Mayer. *Duke Math. J.*, 60(3):599–630, 1990.
- [DM69] P. Deligne and David Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, (36):75–109, 1969.
- [Don81] Ron Donagi. The tetragonal construction. *Bull. Amer. Math. Soc. (N.S.)*, 4(2):181–185, 1981.
- [Don87a] Ron Donagi. Big Schottky. *Invent. Math.*, 89(3):569–599, 1987.
- [Don87b] Ron Donagi. Non-Jacobians in the Schottky loci. *Ann. of Math. (2)*, 126(1):193–217, 1987.
- [Don88] Ron Donagi. The Schottky problem. In *Theory of moduli (Montecatini Terme, 1985)*, volume 1337 of *Lecture Notes in Math.*, pages 84–137. Springer, Berlin, 1988.
- [Don92] Ron Donagi. The fibers of the Prym map. In *Curves, Jacobians, and abelian varieties (Amherst, MA, 1990)*, volume 136 of *Contemp. Math.*, pages 55–125. Amer. Math. Soc., Providence, RI, 1992.
- [DS81] Ron Donagi and Roy Campbell Smith. The structure of the Prym map. *Acta Math.*, 146(1-2):25–102, 1981.
- [Fay73] John D. Fay. *Theta functions on Riemann surfaces*. Lecture Notes in Mathematics, Vol. 352. Springer-Verlag, Berlin, 1973.
- [Fre83] Eberhard Freitag. Die Irreduzibilität der Schottkyrelation (Bemerkung zu einem Satz von J. Igusa). *Arch. Math. (Basel)*, 40(3):255–259, 1983.
- [Gee84] Bert van Geemen. Siegel modular forms vanishing on the moduli space of curves. *Invent. Math.*, 78(2):329–349, 1984.
- [GG86] Bert van Geemen and Gerard van der Geer. Kummer varieties and the moduli spaces of abelian varieties. *Amer. J. Math.*, 108(3):615–641, 1986.
- [GSM07] Samuel Grushevsky and Riccardo Salvati Manni. Singularities of the theta divisor at points of order two. *Int. Math. Res. Not. IMRN*, (15):Art. ID rnm045, 15, 2007.
- [HM98] Joe Harris and Ian Morrison. *Moduli of curves*, volume 187 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998.
- [Igu69] Jun-ichi Igusa. Geometric and analytic methods in the theory of theta-functions. In *Algebraic Geometry (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968)*, pages 241–253. Oxford Univ. Press, London, 1969.
- [Igu81] Jun-ichi Igusa. On the irreducibility of Schottky’s divisor. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 28(3):531–545 (1982), 1981.
- [Iza91] Elham Izadi. *On the moduli space of four dimensional principally polarized abelian varieties*. PhD thesis, Univ. of Utah, 1991.
- [Kri10] Igor Moiseevich Krichever. Characterizing Jacobians via trisecants of the Kummer Variety, arXiv. org: math. *Ann. of Math. (2)*, 2010.
- [MP12] Valeria Ornella Marcucci and Gian Pietro Pirola. Generic Torelli theorem for Prym varieties of ramified coverings. *Compos. Math.*, 148(4):1147–1170, 2012.

- [Mum71] David Mumford. Theta characteristics of an algebraic curve. *Ann. Sci. École Norm. Sup. (4)*, 4:181–192, 1971.
- [Mum74] David Mumford. Prym varieties. I. In *Contributions to analysis (a collection of papers dedicated to Lipman Bers)*, pages 325–350. Academic Press, New York, 1974.
- [Rec74] Sevin Recillas. Jacobians of curves with g_4^1 's are the Prym's of trigonal curves. *Bol. Soc. Mat. Mexicana (2)*, 19(1):9–13, 1974.
- [RF74] Harry E. Rauch and Hershel M. Farkas. *Theta functions with applications to Riemann surfaces*. The Williams & Wilkins Co., Baltimore, Md., 1974.
- [Sch88] F. Schottky. Zur Theorie der Abel'schen Functionen von vier Variablen. *J. Reine und Angew. Math.*, (102):304–352, 1888.
- [Shi86] Takahiro Shiota. Characterization of Jacobian varieties in terms of soliton equations. *Invent. Math.*, 83(2):333–382, 1986.
- [Sho82] V. V. Shokurov. Distinguishing Prymians from Jacobians. *Invent. Math.*, 65(2):209–219, 1981/82.
- [SJ09] F. Schottky and H. Jung. Neue Sätze über Symmetralfunktionen und die Abel'schen Functionen der Riemann'schen Theorie. *S.-B. Berlin Akad. Wiss. (Berlin), Phys. Math. Kl. 1.*, 1909.
- [SM94] Riccardo Salvati Manni. Modular varieties with level 2 theta structure. *Amer. J. Math.*, 116(6):1489–1511, 1994.
- [Wel84] Gerald E. Welters. A criterion for Jacobi varieties. *Ann. of Math. (2)*, 120(3):497–504, 1984.

KAVLI IPMU (WPI), THE UNIVERSITY OF TOKYO, KASHIWA, CHIBA 277-8583, JAPAN

E-mail address: `charles.siegel@ipmu.jp`